



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

		HEDIS®				Non-HEDIS						
		Most recent performance (2017 data)				Most recent performance (2017 data)						
Guideline	Gaps in Care	Measure	Comm	Caid	Care	Measure	Comm	Caid	Care	Comments		
ADHD												
Diagnosis of ADHD for Children and Young Adults		1. Follow-up Care for Children Prescribed ADHD Medication (ADD)										
		a. Initiation Phase and F/U visit	43.38%	43.34%								
		HEDIS® Natl HMO Avg	41.60%	44.60%								
		b. Continuation and Maintenance Phase and F/U Visits	51.05%	52.90%								
Treatment of ADHD for Children and Young Adults		HEDIS® Natl HMO Avg	48.20%	55.00%								
Tools: none												
ASTHMA												
General Principles for the Diagnosis and Management of Asthma	1. Improper use of spacers and inhaling devices 2. Not using lung volume measurements to monitor 3. Inattention to reversing asthma symptoms					1. Preventive visit in year (5-64 years old)	92.84%	89.67%			Recommendations as set forth in the guideline do not directly align with any HEDIS® measure/specifications.	
						2. ED visits with a Primary diagnosis of asthma/1000 asthma members (5-64 years old)	42.6	90.48				
		Tools:										
		NHLBI EPR3:										
		· Classifying Asthma Severity and Initiating Therapy										
		· Assessing Asthma Control and Adjusting Therapy										
		· Stepwise Approach for Managing Asthma Long Term										
		Asthma Initiative of MI: Asthma Action Plans										
		NIH/NHLBI Asthma Action Plan										
		Asthma Guideline Implementation Steps & Tools (GIST)										
How to Use your Asthma Inhaler												
Inhaled Quick Relief and Control Medications (U of M)												
Michigan Asthma Resource Kit (MARK)												
BRONCHITIS												
Management of Uncomplicated Acute Bronchitis in Adults		1. Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB)				33.19%	32.80%					
		HEDIS® Natl HMO Avg				32.00%	33.80%					
		Tools:										
CDC Antibiotic Prescribing and Use in Doctor's Offices												
CDC Preventing and Treating Bronchitis (Patient)												



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments		
		Measure	Most recent performance (2017 data)			Measure	Most recent performance (2017 data)			
			Comm	Caid	Care		Comm	Caid	Care	
CHRONIC KIDNEY DISEASE										
Diagnosis and Management of Adults with Chronic Kidney Disease	GFR calculation necessary					1. Percent members in HEDIS® Diabetes or HTN populations who had a serum creatinine test (necessary for GFR)	94.47%	96.25%	98.29%	No HEDIS® measure Uneven availability of eGFR thru administrative data
		Tools: none								
DEEP VEIN THROMBOSIS										
Outpatient Management of Uncomplicated Deep Venous Thrombosis	1. Pharmacological approaches to outpatient treatment									No measures
		Tools: Anticoagulation Provider Toolkit (MAQI ²) Wells Criteria (DVT scoring system)								
DEPRESSION										
Primary Care Diagnosis and Management of Adults with Depression	1. Pharmacological approaches to treatment	1. Antidepressant Medication Management (AMM)								HEDIS® measures for 84 and 180 days, not "9-12" months after acute symptoms resolve [A]
		a. Effective Acute Phase Treatment	71.96%	57.98%	79.02%					
		HEDIS® Natl HMO Avg	67.80%	53.90%	70.00%					
		b. Effective Continuation Treatment	54.21%	40.92%	62.69%					
		HEDIS® Natl HMO Avg	51.80%	38.60%	55.20%					
		Tools: Edinburgh Postnatal Depression Scale (EPDS) (UCSF) Patient Health Questionnaire (PHQ) for Depression								

*HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments				
		Measure	Most recent performance (2017 data)			Measure	Most recent performance (2017 data)					
			Comm	Caid	Care		Comm	Caid	Care			
DIABETES												
Management of Diabetes Mellitus		1. Comprehensive Diabetes Care (CDC)										
		a. HbA1c Testing	92.40%	88.88%	96.00%							
		HEDIS® Natl HMO Avg	91.20%	87.60%	93.70%							
		b. HbA1c Poor Control (>9.0%)	25.27%	35.52%	14.91%							
		HEDIS® Natl HMO Avg	31.70%	40.50%	25.40%							
		c. HbA1c Control (<8.0%)	63.35%	54.02%	74.19%							
		HEDIS® Natl HMO Avg	57.60%	49.40%	64.40%							
		d. HbA1c Good Control (<7.0%)	n/a	41.32%								
		HEDIS® Natl HMO Avg	n/a	34.50%								
		e. Eye exam, retinal, performed	61.70%	64.67%	80.86%							
		HEDIS® Natl HMO Avg	55.00%	57.20%	71.90%							
		h. Med. Attention for Nephropathy	92.19%	92.29%	96.90%							
		HEDIS® Natl HMO Avg	90.40%	90.10%	95.70%							
		i. Blood Pressure Control (<140/90mmHg)	70.55%	61.93%	73.66%							
		HEDIS® Natl HMO Avg	62.20%	62.70%	67.40%							
		2. Statin Therapy for Patients with Diabetes (SPD)										
		a. Statin Therapy Received	60.74%	65.36%	71.73%							
		HEDIS® Natl HMO Avg	61.50%	61.40%	72.30%							
		b. Statin Therapy Adherence 80%	68.12%	69.17%	74.91%							
		HEDIS® Natl HMO Avg	66.50%	59.00%	74.00%							
		Tools:										
		American Diabetes Association Super Foods for Diabetics										
		Choose My Plate										
		3-Minute Foot Exam for Diabetics										
		Diabetes Checklist (PDF)										
		Diabetes Self-management Education and Support in Type 2 Diabetes (DSME/S)										
		Good Health Club Physician GLs: Prev. & Treatment of Ped Obesity & Diabetes										
		Living with Diabetes An Everyday Guide for You and Your Family										
National Diabetes Education Program Guiding Principles												
Pre-Diabetes:												
AMA/CDC Prevent Diabetes STAT Toolkit												
Combined Diet and Physical Activity Promotion Programs												



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments		
		Measure	Most recent performance (2017 data)			Measure	Most recent performance (2017 data)			
			Comm	Caid	Care		Comm	Caid	Care	
HEART/CARDIOVASCULAR										
Adults with Heart Failure with Reduced Ejection Fraction	1. Underuse ACE & ARBs 2. Treatment for special populations (e.g. African-Americans) 3. Electrolyte monitoring 4. Admission and re-admission avoidance									No HEDIS® measure
Tools: none										
Lipid Screening and Management	1. Pharmacological management with statin therapy	1. Statin Therapy for Patients with Cardiovascular Disease (received therapy and 80% adherence) (SPC)								
		a. Females ages 40-75 received therapy	73.65%	73.88%	76.82%					
		HEDIS® Natl HMO Avg	n/a	n/a	n/a					
		b. Females ages 40-75 80% adherence	69.42%	70.27%	74.87%					
		HEDIS® Natl HMO Avg	n/a	n/a	n/a					
		c. Males ages 21-75 received therapy	83.46%	78.75%	81.67%					
		HEDIS® Natl HMO Avg	n/a	n/a	n/a					
		d. Males ages 21-75 80% adherence	74.08%	71.17%	79.10%					
		HEDIS® Natl HMO Avg	n/a	n/a	n/a					
		e. Total females and males received therapy	81.11%	76.63%	80.06%					
		HEDIS® Natl HMO Avg	80.40%	76.10%	79.00%					
		f. Total females and males 80% adherence	73.07%	70.79%	77.75%					
		HEDIS® Natl HMO Avg	73.50%	63.00%	76.30%					
Tools:										
ACC/AHA ASCVD Risk Estimator Tool										
Choose My Plate										
Statin Dose Intensity and Equivalency Chart										
Medical Management of Adults with Hypertension	1. Pharmacological approaches to treatment	1. Controlling High Blood Pressure (CBP) (single rate reported) -18-59 years of age <140/90 -60-85 years of age w/diabetes <140/90 -60-85 years of age w/o diabetes <150/90	57.57%	58.95%	78.08%	1. Percent members (adult) with any diagnosis of HTN	11.72%	13.56%	44.42%	HEDIS® is ages 18-85 HEDIS® denominator includes (only) members with dx of HTN during first 6 months of the measurement year
		HEDIS® Natl HMO Avg	62.20%	56.90%	70.90%					
Tools:										
FDA Recall of Valsartan Products inc. links to Princeton, Teva, Major Pharm Recall, FDA Updates for health care professionals and patients										
Essential Guide to HTN: Accurate Blood Pressure Readings (AMA)										
CDC Million Hearts - Protocol for Controlling Hypertension in Adults										
DASH Diet										
Hypertension Encounter Form										
Michigan Hypertension Core Curriculum										



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments		
		Measure	Most recent performance (2017 data)		Measure	Most recent performance (2017 data)				
			Comm	Caid	Care		Comm	Caid	Care	
LOW BACK PAIN										
Management of Acute Low Back Pain		1. Use of Imaging Studies for Low Back Pain (LBP)	73.11%	73.31%						
		HEDIS® Natl HMO Avg	76.10%	71.70%						
		Tools:								
		Glossary of Osteopathic Terminology								
		Landmark Healthcare: PT and OT Clinical Practice Guidelines Cervical Segmental Dysfunction								
		Lumbar Spine Segmental Mobility Assessment								
		McKenzie Method Physical Therapy Treatment for Lower Back Pain								
		Spine Segmental Assessment								
OPIOIDS										
Opioids Prescribing in Adults Excluding Palliative and End-of-Life Care										
		Tools:								
		CDC Opioid Prescribing Guidelines Mobile App								
		MDHHS Opioid and Substance Use Disorder Resources								
		Michigan Automated Prescription System (MAPS) inc. Log In link								
		Michigan OPEN: Opioid Prescribing Engagement Network								
		Michigan OPEN: Opioid Prescribing Recommendations for Surgery								
		Michigan Prescription Drug and Opioid Abuse Task Force Report								
		· Task Force Recommendations								
		Opioid Risk Tool (ORT)								
		PEG Scale Assessing Pain Intensity and Interference (Pain, Enjoyment, and General Activity)								
		Screeener and Opioid Assessment for Patients with Pain - Revised (SOAPP-R)								
		Start Talking Form (English, Arabic, Spanish)								
		State of Michigan Opioid Addiction Resources								
ORAL HEALTH										
None										
		Tools:								
		During Pregnancy the Mouth Matters: A Guide to MI Perinatal Oral Health								
		MDHHS Oral Health								
		MI Dental Registry for Healthcare Professionals								
		Recommendations for Preventive Pediatric Health Care: Bright Futures/AAP Periodicity Schedule								
		USPSTS Rec. Statement: Screening for Dental Caries in Children from Birth Through Age 5 Years								
OSTEOARTHRITIS										
Medical Management of Adults with Osteoarthritis	1. Pharmacological approaches to treatment									HEDIS® measure is for DMARD for RA
		Tools:								
		American Arthritis Foundation self-help tools and resources								
OSTEOPOROSIS										
Management and Prevention of Osteoporosis		1. Osteoporosis Management in Women Who Had a Fracture (OMW)		55.14%						HEDIS® measures not sufficient - Medicare and women only - Following fracture, not preventive - Osteoporosis testing (OTO) - from Health Outcomes Survey
		HEDIS® Natl HMO Avg		46.70%						
		Tools:								
		FRAX (WHO Fracture Risk Assessment Tool)								



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

		HEDIS®				Non-HEDIS					
		Most recent performance (2017 data)				Most recent performance (2017 data)					
Guideline	Gaps in Care	Measure	Comm	Caid	Care	Measure	Comm	Caid	Care	Comments	
OVERWEIGHT/OBESITY											
Management of Overweight and Obesity in the Adult	1. Non-surgical interventions for weight loss and weight management	1. Adult BMI Assessment (ABA)	89.50%	94.99%	98.96%						
		HEDIS® Natl HMO Avg	80.30%	84.50%	94.60%						
		Tools:									
		CDC Adult BMI Calculator (20 years and older)									
		Choose My Plate									
Prevention and Identification of Childhood Overweight and Obesity		1. Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)									
		a. BMI Assessment all ages 3-17	87.20%	84.84%							
		HEDIS® Natl HMO Avg	70.30%	72.50%							
		b. Nutrition Counseling all ages 3-17	77.40%	75.79%							
		HEDIS® Natl HMO Avg	64.30%	67.10%							
		c. Coun. For Phys. Activity all ages 3-17	70.39%	68.98%							
		HEDIS® Natl HMO Avg	59.50%	60.60%							
		Tools:									
		AAP Institute for Healthy Childhood Weight: Assessment and Management Algorithm									
		AAP Rx for Healthy Active Living									
CDC BMI Calculator for Children and Teens 2-19 years											
CDC Clinical Growth Charts (Children & Adolescents)											
Good Health Club Physician GLs: Prev. & Treatment of Ped Obesity & Diabetes											
Treatment of Childhood Overweight and Obesity											
PHARYNGITIS											
Acute Pharyngitis in Children 2-18 years	1. Inappropriate use of antibiotics	1. Appropriate testing for children with pharyngitis (CWP)	86.56%	78.85%							
		HEDIS® Natl HMO Avg	86.80%	78.30%							
		Tools:									
		CDC Antibiotic Prescribing and Use in Dotor's Offices: Pediatric Treatment Recommendations									



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

		HEDIS®				Non-HEDIS					
		Most recent performance (2017 data)				Most recent performance (2017 data)					
Guideline	Gaps in Care	Measure	Comm	Caid	Care	Measure	Comm	Caid	Care	Comments	
PREGNANCY PREVENTION											
Prevention of Unintended Pregnancy in Adults 18 Years and Older	1. Interventions to decrease risk of unintended pregnancy									No HEDIS® measure No meaningful measures from administrative data	
		Tools:									
		CDC Planning for Pregnancy, Preconception Health									
		CDC Steps to a Healthier me and baby-to-be! (patient)									
		CDC Summary Chart of US Medical Eligibility Criteria for Contraceptive Use									
Prevention of Pregnancy in Adolescents 12-17 years	1. Interventions to decrease risk of unintended pregnancy in adolescents										
		Tools:									
		CDC Reproductive Health: Teen Pregnancy									
		MDCH Teen Pregnancy Prevention Initiative									
		Michigan Minor Consent Laws									
Provider Tool for Dialogue											
PRENATAL AND POSTPARTUM CARE											
Routine Prenatal and Postnatal Care		1. Prenatal and Postpartum Care (PPC)								HEDIS® frequency of on-going prenatal care (FPC) is Medicaid only. Differences exist among health plans in the use of hybrid method, or report based on administrative data only, thereby foregoing "lift" and reporting lower rates.	
		a. Prenatal: % deliveries that received a prenatal care visit in 1st trimester or within 42 days after enrollment	81.62%	80.82%							
		HEDIS® Natl HMO Avg	84.20%	81.10%							
		b. Postpartum: % deliveries that had postpartum visit on or between 21-56 days after delivery	79.24%	67.82%							
		HEDIS® Natl HMO Avg	74.90%	64.40%							
		Tools:									
		CDC: Pregnancy Education and Resources									
Deliveries Before 39 Weeks (ACOG)											
- March of Dimes: Pregnancy Infographic 39 Weeks											
March of Dimes: Birth Plan for Mom											
Preconception Screening & Counseling Checklist (March of Dimes)											
Women's Health, Pregnancy: Before, During and After											



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments			
		Measure	Most recent performance (2017 data)		Measure	Most recent performance (2017 data)					
			Comm	Caid	Care		Comm	Caid	Care		
PREVENTIVE SERVICES											
ADULTS											
Adult Preventive Services (Ages 18-49)		1. Colorectal Cancer Screening (COL)	70.27%		79.43%					ISSUES: - Specs differ regarding frequency of preventive services Commercial every three years Medicare/Medicaid annually - Age stratifications at 20-44, 45-64, 65+	
		HEDIS® Natl HMO Avg	63.00%		69.60%						
		2. Breast Cancer Screening (BCS)	74.43%	62.44%	80.55%						
		HEDIS® Natl HMO Avg	72.70%	58.30%	72.50%						
		3. Cervical Cancer Screening (CCS)	75.37%	67.14%							
		HEDIS® Natl HMO Avg	74.30%	59.40%							
		4. Chlamydia Screening in Women (ages 21-24) (CHL)	58.23%	70.87%							
		HEDIS® Natl HMO Avg	54.00%	63.70%							
		5. Flu Vaccinations for Adults ages 18-64 (CAHPS) (FVA)	51.67%	39.06%							
		HEDIS® Natl HMO Avg	50.50%	39.60%							
Adult Preventive Services (Age ≥ 50)		6. Flu Vaccinations for Older Adults 65+ (CAHPS) (FVO)			70.28%						
		HEDIS® Natl HMO Avg			72.10%						
		7. Non-recommended PSA-based Screening in Older Men (age 70+) (PSA)			36.51%						
		HEDIS® Natl HMO Avg			31.00%						
		8. Pneumococcal Vaccination Status for Older Adults (PNU)			69.56%						
		HEDIS® Natl HMO Avg			71.30%						
		Tools:									
		ePSS Electronic Preventive Services Selector (AHRQ)									
Adult Immunization Practice Standards											
Everybody Gets Screened Infographic											
National Colorectal Cancer Roundtable Resource Center											
Michigan Breast Density Notification (women)											
Project Accessibility USA (for women)											
Women's Health: Before you get Pregnant											
Advance Care Planning										Care for Older Adults (COA) Four components: Advance Care Planning doc. Medication review Functional status assessment Pain assessment	
Tools: none											



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments				
		Measure	Most recent performance (2017 data)		Measure	Most recent performance (2017 data)						
			Comm	Caid	Care		Comm	Caid	Care			
ADOLESCENTS AND CHILDREN												
Adolescent Health Risk Behavior Assessment												
		Tools: Bright Futures Guidelines for Health Supervision MQIC Guideline Supplement: Resources and Links Rapid Assessment for Adolescent Preventive Services (RAAPS)										
Routine Preventive Services for Children and Adolescents (Ages 2-21)	1. Parent/Child education and counseling 2. Vaccinations	1. Well-Child visits in the 3rd, 4th, 5th and 6th years of life (W34)	80.44%	76.04%							HEDIS® Well-Child visits 3-6 (W34) omits 2-year olds. Consider HEDIS® Children's and Adolescent's access to primary care providers. NOTES: Annual visits for ages 25 months to 6 years (no stratification break at 3 years). HEDIS® includes up to age 19 - no stratification break at 18.	
		HEDIS® Natl HMO Avg	77.20%	73.00%								
		2. Adolescent Well-Care visits (AWC)	49.02%	58.51%								
		HEDIS® Natl HMO Avg	48.40%	53.00%								
		3. Chlamydia screening in Women (ages 16-20) (CHL)	54.62%	64.78%								
		HEDIS® Natl HMO Avg	43.70%	54.20%								
		4. Non-recommended Cervical Cancer Screening in Adolescent Females	1.49%	2.07%								
		HEDIS® Natl HMO Avg	1.50%	1.50%								
		5. Immunizations for Adolescents (IMA)										
		a. Meningococcal	90.86%	89.06%								
		HEDIS® Natl HMO Avg	81.10%	79.60%								
		b. Tdap	91.45%	87.15%								
		HEDIS® Natl HMO Avg	88.00%	86.20%								
		c. Combo 1	89.34%	85.41%								
		HEDIS® Natl HMO Avg	79.70%	77.60%								
		d. Combo 2	27.21%	36.50%								
		HEDIS® Natl HMO Avg	24.00%	32.70%								
		e. Human Papillomavirus Vaccine	27.90%	38.24%								
		HEDIS® Natl HMO Avg	25.70%	35.30%								
				Tools: AAP Bright Futures: Recommendations for Preventive Pediatric Health Care Periodicity Schedule Alliance for Immunization in Michigan Toolkit CDC Childhood Vaccine Quiz for Parents CDC Vaccines and Immunizations: Parents, What You Need to Know CDC Video: Wash Your Hands I Vaccinate MDHHS Adolescent Immunization Website								



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments	
		Measure	Comm	Caid	Care	Measure	Comm		Caid
Routine Preventive Services for Infants and Children (Birth-24 months)	1. Parental education and counseling 2. Vaccinations	1. Childhood Immunization Status (CIS)							
		a. DTaP	86.97%	78.79%					
			HEDIS® Natl HMO Avg	85.30%	76.30%				
		b. IPV	92.22%	90.35%					
			HEDIS® Natl HMO Avg	90.60%	88.20%				
		c. MMR	91.21%	90.63%					
			HEDIS® Natl HMO Avg	92.50%	88.80%				
		d. Hib	92.21%	88.23%					
			HEDIS® Natl HMO Avg	91.60%	87.70%				
		e. HepB	91.11%	92.31%					
			HEDIS® Natl HMO Avg	86.30%	88.20%				
		f. VZV	90.70%	90.06%					
			HEDIS® Natl HMO Avg	92.00%	88.60%				
		g. Pneumococcal conjugate	87.45%	77.93%					
			HEDIS® Natl HMO Avg	85.70%	76.80%				
		h. HepA	86.04%	86.37%					
			HEDIS® Natl HMO Avg	87.80%	84.30%				
		i. Rotavirus	82.54%	71.02%					
			HEDIS® Natl HMO Avg	81.80%	70.10%				
		j. Influenza	62.30%	44.70%					
			HEDIS® Natl HMO Avg	66.10%	47.40%				
		k. Combo 2	82.80%	77.01%					
			HEDIS® Natl HMO Avg	78.60%	72.90%				
		l. Combo 3	81.23%	72.84%					
			HEDIS® Natl HMO Avg	76.80%	69.40%				
		s. Combo 10	54.31%	36.09%					
			HEDIS® Natl HMO Avg	53.40%	35.40%				
		2. Well-Child visits in the first 15 months of life 6+ visits (W15)	84.83%	72.17%					
			HEDIS® Natl HMO Avg	78.20%	64.10%				
		3. Lead Screening in Children (LSC) [Medicaid only]		80.24%					
			HEDIS® Natl HMO Avg	68.90%					
Tools:									
AAP Bright Futures: Recommendations for Preventive Pediatric Health Care Periodicity Schedule									
Alliance for Immunization in Michigan Toolkit									
CDC Childhood Vaccine Quiz for Parents									
CDC Vaccines and Immunizations: Parents, What You Need to Know									
CDC Video: Wash Your Hands									
I Vaccinate									
WHO Child Growth Charts (Birth to 24 months)									



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

		HEDIS®				Non-HEDIS				
		Most recent performance (2017 data)				Most recent performance (2017 data)				
Guideline	Gaps in Care	Measure	Comm	Caid	Care	Measure	Comm	Caid	Care	Comments
SEDATION USE, OFFICE										
In Office Use of Sedation										
Tools: none										
SUBSTANCE USE										
Screening, Diagnosis and Referral for Substance Use Disorders	1. Assessment and identification of substance use disorders	1. Initiation and Engagement of Alcohol and Other Drug (AOD) Dependence Treatment (IET)								HEDIS 2018 1st yr reporting
		a. Initiation of AOD treatment ALCOHOL	35.29%		47.54%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		b. Engagement of AOD treatment ALCOHOL	12.10%		5.82%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		a. Initiation of AOD treatment OPIOID	39.35%		37.47%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		b. Engagement of AOD treatment OPIOID	17.58%		6.25%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		a. Initiation of AOD treatment OTHER	36.12%		42.72%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		b. Engagement of AOD treatment OTHER	9.52%		3.05%					
		HEDIS® Natl HMO Avg	n/a		n/a					
		a. Initiation of AOD treatment ALL	34.46%		43.14%					
		HEDIS® Natl HMO Avg	37.10%		32.60%					
		b. Engagement of AOD treatment ALL	11.91%		5.40%					
		HEDIS® Natl HMO Avg	12.30%		3.80%					
Tools:										
CRAFTT (adolescents)										
MDHHS Substance Use, Problem Gaming, or Mental Health Contact Information										
Michigan Automated Prescription System (MAPS)										
National Institute on Alcohol Abuse and Alcoholism Treatment Navigator										
SAMHSA - HRSA Center for Integrated Health Solutions (screening tools)										
SAMHSA - Michigan Buprenorphine Locator										
TWEAK - pregnant women										



MQIC GUIDELINES, MEASURES and TOOLS SUMMARY
MQIC Community Results - HEDIS® 2018 (Measurement Year 2017)

Guideline	Gaps in Care	HEDIS®			Non-HEDIS			Comments			
		Measure	Most recent performance (2017 data)			Measure	Most recent performance (2017 data)				
			Comm	Caid	Care		Comm	Caid	Care		
TOBACCO USE											
Tobacco Control	1. Identification of tobacco use and exposure 2. Interventions to decrease tobacco use	1. Medical Assistance with Smoking Cessation (MSC) NOTE: CAHPS measure - Adults - 18+				1. Percent Current Smokers (calculated in Quality Compass) - Age 18+	11.13%	40.26%	n/a	Guideline: Age 12+ Project to total number of current adult smokers (18+) based on % current smokers and number of adults 18+ (available in HEDIS®)	
		a. Advising Smokers to Quit	79.78%	81.64%	68.99%						
		HEDIS® Natl HMO Avg	75.90%	77.00%	86.20%						
		b. Discussing Smoking Cessation Medications	53.33%	58.87%							
		HEDIS® Natl HMO Avg	52.50%	51.50%							
		c. Discussion Smoking Cessation Strategies	44.49%	47.75%							
		HEDIS® Natl HMO Avg	45.70%	45.40%							
		Tools:									
		CDC Million Hearts: Protocol for Identifying and Treating Patients Who Use Tobacco									
		CDC Million Hearts: Tobacco Cessation Protocol Implementation Guidance									
Guide Your Patients to a Smoke-Free Future (PDF) (Canada)											
MI Steps Up "Don't Smoke" (comprehensive patient and physician materials)											
MI Cancer Consortium Tobacco Cessation Provider Toolkit											
MI Medicaid Tobacco Cessation Benefits											

OTHER TOOLS

Michigan Department of Health and Human Services	
Access to Primary Care for People with Disabilities	
American Disability Act: Access to Medical Care for Individuals with Mobility Disabilities	
I Vaccinate	
Michigan Health and Wellness 4x4 Plan	
Oral Health	
Reporting Requirements:	
Michigan Care Improvement Registry	
Reportable Diseases in Michigan	
Requirements for Reporting Abuse and Neglect	

KEY

No measure or no data	
Guideline does not apply	
Outperformed HEDIS® 2018 Natl Average	
Underperformed HEDIS® 2018 Natl Average	
HEDIS® 2018 Natl Avg not avail	n/a

MQIC RESULTS (PREVIOUS YEAR COMPARISON)

Decline from previous year result
Increase from previous year result
Not previously reported no comparison